UCI - 90 - 33
June 1990

受入
90-7-278
高工研図書室

Critical Properties of Two-Dimensional Simplicial Quantum Gravity

Mark Gross [1] *and Herbert W. Hamber* [2]

[1] Department of Physics and Astronomy
California State University at Long Beach, Long Beach, Ca 90840 USA

[2] Department of Physics
University of California at Irvine, Irvine, Ca 92717 USA

ABSTRACT

Simplicial higher derivative quantum gravity is investigated in two dimensions for a manifold of toroidal topology. The manifold is dynamically triangulated using Regge's formulation of gravity, with continuously varying edge lengths and fixed coordination number. Critical exponents are estimated by computer simulation on lattices with up to 786432 edges, and compared to the continuum conformal field theory results for central charge zero (pure gravity), one half (Ising model coupled to gravity), one and two (massless scalar field coupled to gravity). The dependence of critical properties on the coefficient of the curvature squared term and the gravitational functional measure is investigated, suggesting universal critical behaviour at least within a certain class of measures. In the case of pure gravity, we have computed the string susceptibility exponent for both the torus and the sphere, and our estimates agree with the exact result of KPZ. The fluctuations in the area density are consistent with the behavior expected for a massless scalar field, the Liouville mode. In the case of gravity coupled to a massless scalar field, we have computed what corresponds to the fractal dimension of the surface, and found it to be infinite. The critical exponents associated with the Ising model coupled to gravity on a torus are found to be the same as for the Ising model in flat space.

1. INTRODUCTION

Understanding properties of two-dimensional lattice quantum gravity is likely to be an essential step on the way to formulating a lattice theory of four-dimensional quantum gravity. In addition, two-dimensional quantum gravity is closely related to the theory of random surfaces embedded in D-dimensional flat space, as well as to the theory of bosonic strings.

We will concentrate here on the simplicial formulation of quantum gravity developed by Regge[1-12], but it is known that there are other possible approaches such as the random triangulation approach for two-dimensional surfaces[13-17], as well as the hypercubic lattice formulation[18,19]. It would not be too surprising if these three approaches are someday shown to be equivalent, since after all they correspond to different discretizations of the same original theory. On the other hand their quantum continuum limit could in principle exhibit rather different properties, and in particular there is certainly no general proof yet of the recovery of general coordinate invariance.

One of the advantages of the simplicial Regge calculus approach lies in the fact that it can be formulated in any space-time dimension (including the physically relevant case of four dimensions), and in the fact that it can be shown to be classically equivalent to general relativity, as a consequence of several more or less rigorous convergence proofs for (arbitrarily) triangulated smooth manifolds[2-5,7,9,12]. In addition, the correspondence between the lattice and continuum quantities is clear, and therefore the interpretation of the terms in the action as well as the identification and separation of, for example, the measure contribution is unambiguous[6]. For a review of Regge gravity, and a more complete list of early references, we refer the reader to refs. [6].

Approximate general coordinate (or re-parametrization) invariance corresponds to variations in the edge lengths that leave the geometry of the underlying manifold unchanged. For the limiting case of flat space there are clearly infinitely many edge length assignments which reproduce equally well the underlying manifold. This situation can be contrasted to the case of a triangulation with fixed edge lengths and a varying incidence matrix, for which there is no re-parametrization invariance (except for the trivial one, corresponding to a re-labelling of the lattice vertices). As one moves away from flat space though, different edge lengths assignments will in general correspond to physically inequivalent manifolds.

In continuum gravity the fundamental degrees of freedom are represented by the metric field $g_{\mu\nu}(x)$. In a two-dimensional piecewise linear space the elementary building blocks are triangles, and the relative position of points on the lattice is therefore completely specified by the incidence matrix and the edge lengths, which in turn induces a metric structure on the piecewise linear space. In order to obtain non-degenerate simplicial complexes, the edge lengths have to obey triangle inequalities, which ensure that the triangles areas are positive. General coordinate transformations correspond (at least approximately) to variations of the edge lengths, as well as appropriate modifications of the incidence matrix. But since in general different complexes will correspond to physically distinct manifolds, one expects classically general coordinate invariance to be recovered only in the continuum limit, where a continuous smooth manifold can be covered by many different almost geometrically equivalent triangulations. In the quantum theory the hope is to find a non-trivial fixed point where general coordinate invariance is recovered.

A detailed description of the construction of the action for simplicial lattice gravity without and with matter fields can be found in the literature [5,6,9,11], and therefore only a brief summary will be presented here. Ref. [9] addressed the specific case of pure gravity in two dimensions, and discusses some analytic and numerical results related to the phase diagram of pure gravity. We recall the geometric correspondence between continuum and lattice quantities in two dimensions[5,9]

$$\begin{aligned} \int d^2x\,\sqrt{g(x)} &\rightarrow \sum_i A_i \\ \int d^2x\,\sqrt{g(x)}\,R(x) &\rightarrow 2\sum_i \delta_i \\ \int d^2x\,\sqrt{g(x)}\,R^2(x) &\rightarrow 4\sum_i \delta_i^2/A_i \end{aligned} \tag{1.1}$$

where δ_i is the deficit angle at the vertex i

$$\delta_i = 2\pi - \sum_{\substack{\text{triangles} \\ \text{meeting on } i}} \theta_d \tag{1.2}$$

and θ_d is the dihedral angle associated with the triangle at that vertex. A_i is the area associated with the site i [5,9], which is not unique since the lattice can be subdivided in more than one way, for example using a dual lattice or a baricentric subdivision. Given reasonable geometric and positivity properties, universality is expected to lead to the same results in the continuum. Note that in the simplicial lattice formulation, as in the continuum, the local curvature $R(x) \sim 2\delta_i/A_i$ is a continuous function of the relevant edge lengths and can take on any real value, positive or negative. In two dimension the discrete analogue of the Gauss-Bonnet theorem holds

$$\sum_i \delta_i = 2\pi\chi \tag{1.3}$$

where χ is the Euler characteristic of the surface (two minus twice the number of handles). In this paper we will consider only simplicial complexes topologically equivalent to the torus and the sphere.

The guiding principle in constructing physical quantities in simplicial gravity is that they should have geometric significance. It will distinguish objects which are lattice structure independent for a given physical manifold (at least for sufficiently smooth manifolds in some continuum limit) from other functions of the edge lengths which have no particular geometric meaning, and whose limiting values will therefore depend on the specific way in which the triangulation is refined. The Euler characteristic in two dimensions, expressed as a function of the edge lengths, is a clear and illustrative example of what is meant by this statement. Another clear example is the total area of the simplicial complex: if it is defined as the sum of the triangle areas (where these are very specific functions of the edge lengths), then as the triangulation is refined its limit is well defined, and agrees with the continuum definition of what is meant by the total physical area of the manifold.

Having made some general considerations regarding the action for gravity, let us now turn to the issue of the measure. The form of the measure for the $g_{\mu\nu}$ fields in continuum gravity appears not to be unique, and the topic has been discussed recurrently in the literature [20-23]. The reason for the ambiguities appears to be a lack of a clear definition of what is meant by $\prod_x$ in the functional measure. Thus it is expected that the ambiguities will persist in *any* lattice formulation of quantum gravity, unless an exact lattice invariance is found, which then uniquely selects one privileged measure. However, as will be discussed below, different measures differ by the power of $\sqrt{g}$ in the prefactor, which corresponds to some product of volume factors on the lattice.

One popular (pure) gravitational measure in the continuum is the Misner scale-invariant measure[20], which in d space-time dimensions takes the form

$$d\mu[g] = \prod_x g^{-(d+1)/2} \prod_{\mu\geq\nu} dg_{\mu\nu} \tag{1.4}$$

The above measure is unique if the product in eq. (1.4) is interpreted over 'physical' points, and invariance is imposed at one and the same 'physical' point. On the other hand, if one introduces a super-metric over metric deformations, then another measure arises naturally for pure gravity[21]. Considering the simplest local form for the norm-squared of the metric deformation

$$\begin{aligned} ||\delta g||^2 &= \frac{1}{2}\int d^dx\sqrt{g}\left[g^{\mu\alpha}g^{\nu\beta} + g^{\mu\beta}g^{\nu\alpha} + \lambda g^{\mu\nu}g^{\alpha\beta}\right]\delta g_{\mu\nu}\delta g_{\alpha\beta} \quad (\lambda \neq -\frac{2}{d}) \\ &\equiv \int d^dx\sqrt{g}\, G^{\mu\nu,\alpha\beta}\,\delta g_{\mu\nu}\delta g_{\alpha\beta} \end{aligned} \tag{1.5}$$

then, according to DeWitt, the functional measure is given by

$$d\mu[g] = \prod_x \left[\det G^{\mu\nu,\alpha\beta}\right]^{1/2} \prod_{\mu\geq\nu} dg_{\mu\nu} \tag{1.6}$$

where the determinant of the super-metric G is given by

$$\left[\det G^{\mu\nu,\alpha\beta} \right]^{1/2} = (-1)^{d-1}(1+\frac{d\lambda}{2})\, g^{(d-4)(d+1)/8} \tag{1.7}$$

If matter fields are present, then the gravitational measure has to be further modified. Other forms of the measure for the gravitational field have also been suggested, inspired by the canonical quantization approach to gravity[22].

On the simplicial lattice the edge lengths are the elementary degrees of freedom which uniquely specify the geometry for a given incidence matrix, and over which one should perform the functional integral[5,6,8,9]. This is supported by the fact that the induced metric on a simplex i with edges $l_{i,i+\mu}$ is given by $g_{\mu\nu}(i) = \frac{1}{2}\left[l^2_{i,i+\mu} + l^2_{i,i+\nu} - l^2_{i+\mu,i+\nu} \right]$. In addition, one might want to sum over complexes with varying local coordination number. But it would seem that any smooth curved manifold can be arbitrarily well approximated by a simplicial complex with fixed coordination number by adjusting the edge lengths and refining the grid, thus presumably saturating the functional measure. This would suggest that the last step is then perhaps redundant, at least within our formalism.

One can argue that the edge lengths, being invariant quantities, are not referred to any specific coordinate systems. On the other hand they provide for an explicit coordinatization of the manifold, once the incidence matrix is specified as well. It is clear from looking at the example of flat space that there can be an infinite number of edge length assignments that correspond to the same physical manifold. Therefore in the continuum limit the edge lengths cannot really be considered as invariants under some (approximate) lattice diffeomorphism group.

Previously[5,8,9,6] we have employed the measure

$$\int d\mu_\epsilon[l] = \prod_{\text{edges ij}} \int_0^\infty \frac{dl^2_{ij}}{l^2_{ij}}\, F_\epsilon[l] \tag{1.8}$$

where $F_\epsilon[l]$ is a function of the edge lengths with the property that it is equal to one whenever the triangle inequalities are satisfied, and zero otherwise. A parameter ϵ can be introduced as an ultraviolet cutoff at small edge lengths: the function $F_\epsilon[l]$ is then chosen to be zero if any of the edges are equal or less than ϵ; in the following we will take $\epsilon = 0$. The above measure is correct in the weak field limit[2,9], where all continuum measures agree as well.

It is of interest though to explore the sensitivity of the results to the type of gravitational measure employed. Another class of pure gravity measures is obtained by considering the 'volume associated with an edge' V_{ij}, and writing in two dimensions

$$\int d\mu_\epsilon[l] = \prod_{\text{edges ij}} \int_0^\infty V_{ij}^{2\sigma}\, dl^2_{ij}\, F_\epsilon[l] \tag{1.9}$$

with $\sigma = -1/2$ for the lattice analog of the Misner measure, and $\sigma = -1/4$ for a lattice analogue of the DeWitt measure. Note that the 'Misner' and dl/l measure share the property of being scale invariant. Most of our results will be based on the measure dl/l. But we will explore the properties of these two other lattice measures as well, and shall return to the issue of the measure when we discuss the coupling of gravity to a scalar field, and how the results depend on the form of the measure and on σ. Our results suggest that different measures, within a certain universality class, will give the same results for infrared sensitive quantities, like correlation functions at large distances and critical exponents. We believe though that the lattice path integral might not be meaningful for certain values of σ. We have found in particular that if σ is too negative, then the measure factors tend to favor configurations of triangles which are long and thin, with a small area and a large perimeter. In this case the lattice tends to degenerate into an almost one-dimensional manifold, a situation far from the desired continuum limit, and which one would like to avoid.

In two dimensions a measure for gravity has been given by Polyakov[24,25-27], following the DeWitt approach. In pure two-dimensional gravity, and for vanishing renormalized cosmological constant $\lambda \to \lambda_c$, one can write for the path integral

$$\int d\mu[g]\, \exp\left(-I_G\right) = \int d\mu[\tilde{g}]\, \Delta_{FP}[\tilde{g}] \int [d\varphi]\, \exp\left(-I_G - 26 I_L\right) \tag{1.10}$$

with the Liouville action contribution I_L arising from the conformal anomaly

$$I_L = \frac{1}{96\pi} \int d^2x \sqrt{\tilde{g}} \, (\tilde{g}^{\mu\nu} \partial_\mu \varphi \, \partial_\nu \varphi + 2\bar{R} \, \varphi) \tag{1.11}$$

Here $g_{\mu\nu}(x) = \tilde{g}_{\mu\nu}(x) e^{\varphi(x)}$, and $\tilde{g}_{\mu\nu}(x)$ is some reference metric. Thus for vanishing renormalized cosmological constant one expects the relevant gravitational degrees of freedom to be represented by a free scalar field. On the lattice the conformal factors correspond to local area fluctuations, $e^{\varphi(x)} \approx V(x)/V_0$. In the following sections we will discuss some of the properties of the density fluctuation field φ in simplicial gravity, which are in agreement with the continuum predictions.

Let us now turn to presentation of our results. In the following section we will discuss properties of pure gravity, including the critical exponent γ_χ for the torus and the sphere, and the Liouville field susceptibility. Then we will present results for a model of gravity coupled to a D-component massless scalar field, and make contact with models of random surfaces (here without extrinsic curvature terms). We will consider the cases $D = 1$ and 2, as well as the case $D = 0$, which corresponds an absence of feedback of the matter fields on the geometry. Finally we will compute the critical exponents for an Ising model coupled to gravity on a torus. The last section presents our conclusions.

2. PURE GRAVITY

We will begin by considering a higher derivative lattice action for pure gravity in the form

$$I = \sum_i \left[\lambda \, A_i - k \, \delta_i + a \, \frac{\delta_i^2}{A_i} \right] \tag{2.1}$$

For a detailed description of the construction of curvature terms on the simplicial lattice we refer to ref. [5]. In the following we will adopt a baricentric subdivision for assigning area elements to the lattice sites and edges. One could use a dual lattice subdivision as well, but it would lack some desirable positivity properties for the dual areas in some cases[5]. In the limit of small fluctuations around a smooth background, the above lattice action can be shown to correspond to the continuum action

$$I = \int d^2x \sqrt{g} \left[\lambda - \frac{k}{2} \, R + \frac{a}{4} \, R^2 \right] \tag{2.2}$$

For a manifold of fixed topology the term proportional to k can be dropped. The higher derivative term proportional to a can be useful in controlling the fluctuations in the intrinsic local curvature, although it is expected to be irrelevant as far as critical properties are concerned. While it prevents the appearance of conical singularities where the Gaussian curvature might become very large, it does not prevent 'folding' singularities, corresponding to singular structures in the manifold which appear in embedding space only (to control the latter an extrinsic curvature term seems to be required). For $a \to \infty$ the manifold approaches a flat limit, whereas for $a \to 0$ local fluctuations in the curvature become more pronounced. In ref. [9] it was shown that no sensible ground state exists for $a < 0$ (unless there are additional terms in the action), and in the following we will therefore only discuss the case $a \geq 0$.

Classically the continuum equations of motion lead to a constant curvature solution $R_0(x) = \pm\sqrt{4\lambda/a}$, (there being no real solution for $\lambda < 0$). On the torus on the other hand the only consistent classical solution is $R_0(x) \equiv 0$, an identity which remains true for the average when quantum fluctuations are included, as a consequence of the Gauss-Bonnet theorem.

In ref. [9] the canonical ensemble (with fixed bare cosmological constant λ) was considered. On the other hand, in order to compare with the exact results of KPZ[26], it is useful to consider an ensemble where the total area A is kept fixed instead. In such an ensemble the limit $A \to \infty$ corresponds to $\lambda \to \lambda_c$ in the canonical ensemble. One approaches the limit $A \to \infty$ by letting the number of sites $N \to \infty$ for fixed elementary triangle areas (to approach the continuum limit with a fixed total physical area, the elementary

triangle areas would have to be scaled to zero as N is increased, which is not what we shall do here). We will therefore consider the lattice analogue of

$$Z\,[A] = \int d\mu[g]\;\delta(\int \sqrt{g} - A)\;\exp\,(-I[g]) \tag{2.3}$$

which for large area is expected to behave asymptotically as

$$Z\,[A] \underset{A\to\infty}{\sim} A^{-3+\gamma_\chi}\;\exp\,(-(\lambda-\lambda_0)A) \tag{2.4}$$

where λ_0 affects the renormalization of the cosmological constant. The exponent $\gamma_\chi = \chi(\gamma-2)/2+2$, with $\gamma = \frac{1}{12}(D-1-\sqrt{(D-1)(D-25)})$, is the 'string susceptibility' exponent[26-28]. Note that the exponent γ_χ is not truly universal, since it depends on the boundary conditions through χ. Pure gravity without matter fields then corresponds to the case $D=0$, and in particular on the torus one has the prediction $\gamma_\chi = 2$, independent of D.

It is easy to see that the exponent γ_χ can be related to a finite size correction. By doing an infinitesimal scale transformation on $Z[A]$, with the action given by eq. (2.2), one obtains the identity

$$\frac{\partial \ln Z[A]}{\partial A} = -\frac{1}{A} + \frac{a}{4}\,\frac{<\int\sqrt{g}\,R^2>_A}{A} + \lambda_0' - \lambda \tag{2.5}$$

where λ_0' depends, among other things, on the specific form of the measure. Using eq. (2.4) for $Z[A]$ one then has

$$\frac{a}{4}\,\frac{<\int\sqrt{g}\,R^2>_A}{A} \underset{A\to\infty}{\sim} \text{const} - \frac{2-\gamma_\chi}{A} + \cdots \tag{2.6}$$

Thus the critical exponent γ_χ can be obtained by investigating the area dependence of the expectation value of R^2. In particular on the torus one has the prediction that the correction proportional to $1/A$ must have a vanishing coefficient. On the lattice the appropriate quantity to measure is

$$\frac{(1/N)<\sum_i \delta_i^2/A_i>}{(1/N)<\sum_i A_i>} \sim \frac{1}{4}\,\frac{<\int\sqrt{g}\,R^2>_A}{A} \tag{2.7}$$

where N is the total number of sites. Without loss of generality one can choose the total area such that $N = A$.

We have computed the coefficient of the finite size correction to R^2 by investigating both the torus and the sphere, using as a background space a network of unit squares divided into triangles by drawing in parallel sets of diagonals, as shown in Fig. 1. (For the sphere the two poles have to be treated separately due to the presence of a coordinate or incidence matrix singularity). Ideally one would like to use a random lattice[30], but this presents additional computational problems, so we have opted for the moment for the simpler approach of using a regular lattice. On the basis of universality one would expect the results to be independent of the specific lattice structure chosen. Our results indicate that the expected result is clearly obtained for the torus, and with somewhat larger errors for the sphere as well, with a choice of fixed coordination number and varying edge lengths.

In both cases the lattices considered contained from 48 to 12288 edges (corresponding to lattices with 4^2, 5^2, 6^2, 8^2, 10^2, 12^2, 16^2, 32^2 and 64^2 sites). The lengths of the runs varied between 4700k sweeps on the small lattice and 20k sweeps on the largest lattice. The coupling a was set $= 1$, and we used the measure dl/l of eq. (1.8). The results for R^2 versus $1/A$ are shown in Table I, and in Figs. 2 and 3. In the case of the torus the results for the coefficient of the $1/A$ term are quite accurate, and consistent with zero to within a few percent: we obtain from a straight line fit approximately $2-\gamma_\chi = 0.025(7)$ from lattices $8-64$, and $2-\gamma_\chi = 0.047(13)$ for lattices $10-32$, leading to a combined estimate $2-\gamma_\chi = 0.025(22)$. We therefore conclude that for the torus $2-\gamma_\chi$ is very close to, and given our statistical errors almost consistent with, zero.

In Fig. 4 we show as an example the distribution of curvatures δ_i (corresponding to the continuum $R\sqrt{g}$) on a lattice with 262144 sites, with $a = 0.001$. Note that the distribution is a rather smooth function, as is the distribution of edge lengths and triangle areas.

Let us now turn to a discussion of the case of the sphere. On the (almost) regularly triangulated sphere, the results for γ_χ are not quite as accurate as for the torus, since the coordinate singularities introduced at the poles, where many ($\sqrt{N}$) edges meet at one point, introduce a finite size correction proportional to $A^{-3/2}$, which was absent in the case of the torus. This could be avoided by using a different triangulation for the sphere which does not exhibit this feature. Still, by fitting the results for R^2 to $c_0 + c_1/A + c_2/A^{3/2}$ we have obtained $c_0 = 0.423(2)$, $c_1 = -2.75(32)$ and $c_2 = 53.3(2.9)$, and from the coefficient of the $1/A$ term the estimate $\gamma = -0.75 \pm 0.32$, to be compared to the expected KPZ answer of $\gamma = -1/2$. The χ^2 in the fits in general varies between 10 and 200. Our error estimates are obtained by comparing fits that include all or only part of the data points, with proper weighting to take into account the statistical uncertainty in the Monte Carlo data. We have checked that our results are stable when we add an additional term, $+c_4/A^2$, for whose coefficient we find $c_4 = 14.7(15.)$, indeed a relatively small correction in the range of A considered. It is fair to say that in the case of the sphere the results can only be shown to be roughly consistent with the KPZ exact result, although they clearly seem to exclude for example the semiclassical result $\gamma \to -\infty$. The preceding results would suggest a restoration of general coordinate invariance at large distances or low momenta in the lattice theory. In the case of the sphere the results appear to be consistent with the KPZ result, but the errors are quite large. Further tests can be performed by embedding the surface and measuring its extent in embedding space, as will be discussed in the next section.

We have also investigated the critical properties of the area fluctuation or Liouville field $\varphi(x)$, again in the case of pure gravity. We define the discrete analogue of the continuum Liouville field $\varphi(x) = \ln\sqrt{g(x)}$ as $\varphi_i = \ln A_i$, and compute the Liouville-field susceptibility on a finite lattice

$$\chi_\varphi(L) = A\,[\,<\varphi^2> - <\varphi>^2\,] \tag{2.8}$$

with

$$\varphi = \frac{1}{A}\sum_i \ln A_i \tag{2.9}$$

For the measure dl/l at $a = 0$ we find, using finite size scaling on tori of sizes $L = \sqrt{A} = 8 - 128$ (see Table II and Fig. 5),

$$\ln\chi_\varphi(L) \underset{L\to\infty}{\sim} c + (2 - \eta_\varphi)\,\ln L \tag{2.10}$$

with an exponent $2 - \eta_\varphi = 2.08(12)$. This result is consistent with the expected free field behavior of the massless Liouville mode ($\eta_\varphi = 0$), and again suggests a restoration of general coordinate invariance in the quantum theory.

3. GRAVITY COUPLED TO A MASSLESS SCALAR FIELD

Matter fields are introduced in a straightforward way. Consider a D-component scalar field ϕ_i^a, $a = 1, ..., D$. Define the scalar fields at the vertices of the triangles, and introduce finite lattice differences defined in the usual way[11,6]

$$(\Delta_\mu \phi^a)_i = \frac{\phi^a_{i+\mu} - \phi^a_i}{l_{i,i+\mu}} \tag{3.1}$$

The index μ labels the possible directions in which one can move from a point in a given triangle, and $l_{i,i+\mu}$ is the length of the edge connecting the two points. As an action we choose

$$I\,[\phi] = \frac{1}{2}\sum_{\textbf{edges ij}} V_{ij}\left(\frac{\phi^a_i - \phi^a_j}{l_{ij}}\right)^2 \tag{3.2}$$

where V_{ij} is the volume associated with the edge ij, via a baricentric subdivision,

$$V_{ij} = \sum_{\textbf{triangles t}\,\supset\,\textbf{ij}} \frac{1}{3}\,V_t \tag{3.3}$$

The above lattice action then corresponds to the continuum expression

$$\frac{1}{2}\int d^2x\,\sqrt{g}\,g^{\mu\nu}\,\partial_\mu\phi^a\,\partial_\nu\phi^a \tag{3.4}$$

Other forms for the scalar field action have been suggested[11], and we expect them to be equivalent in some continuum limit. The discretized partition function is then given by

$$Z=\int d\mu[l]\,d\mu[\phi]\,\exp\left(-I[l]-I[\phi]\right) \tag{3.5}$$

Let us now consider what the appropriate lattice measure should be. For a D-component scalar field the continuum scalar field measure[21]

$$d\mu[\phi]=\prod_x g^{D/4}(x)\prod_a d\phi^a(x) \tag{3.6}$$

leads to a combined continuum measure for the gravitational and scalar degrees of freedom

$$d\mu[g]=\prod_x\prod_{\mu\geq\nu} g^\sigma(x)\,dg_{\mu\nu}(x)\prod_x\prod_a d\phi^a(x) \tag{3.7}$$

with $\sigma=(D-1)/4$ for the DeWitt measure, and $\sigma=(D-2)/4$ for the Misner measure. Their discretized form then reads

$$\int d\mu_\epsilon[l]\int d\mu[\phi]=\prod_{\text{edges ij}}\int_0^\infty V_{ij}^{2\sigma}\,dl_{ij}^2\,F_\epsilon[l]\prod_{\text{sites i}}\prod_a\int_{-\infty}^{\infty}d\phi_i^a \tag{3.8}$$

Again we have also considered the simpler dl/l-type measure[8,9,6]

$$\int d\mu_\epsilon[l]\int d\mu[\phi]=\prod_{\text{edges ij}}\int_0^\infty\frac{dl_{ij}^2}{l_{ij}^2}\,F_\epsilon[l]\prod_{\text{sites i}}\prod_a\int_{-\infty}^{\infty}d\phi_i^a \tag{3.9}$$

All the above forms of the lattice measure can be recast in the form

$$\prod_{\text{edges ij}}\int_0^\infty\frac{dl_{ij}^2}{l_{ij}^2}\,F_\epsilon[l]\prod_{\text{sites i}}\prod_a\int_{-\infty}^{\infty}d\phi_i^a\,\exp\Big(\sum_{ij}\ln\left[\,l_{ij}^{2\alpha}V_{ij}^{2\sigma}\,\right]\Big) \tag{3.10}$$

with $\alpha=\sigma=0$ for the dl/l measure, and $\alpha=1$ and σ equal to the values mentioned above for the lattice analogues of the DeWitt and Misner measure. Most of our simulation results to be presented later refer to the dl/l measure. But we will show results and give arguments which suggest that all three measures lead, in the cases we have considered, to the same critical properties.

In order to study the properties of the scalar field coupled to gravity, and attempt to compare with related work, we have measured the discrete analog of the coordinate invariant quantity

$$<\phi^2>=\frac{1}{D}\,\frac{<\int\sqrt{g}\,(\phi^a-\bar{\phi}^a)^2>}{<\int\sqrt{g}>} \tag{3.11}$$

with

$$\bar{\phi}^a=\frac{\int\sqrt{g}\,\phi^a}{\int\sqrt{g}} \tag{3.12}$$

Since we are working in an ensemble in which the total area is fixed, and equal to the number of sites, we have $\int\sqrt{g}=\sum_i A_i=N=A$. On the lattice we measure therefore

$$<\phi^2>=\frac{1}{DN}<\sum_i A_i\,(\phi_i^a-\bar{\phi}^a)^2> \tag{3.13}$$

with

$$\bar{\phi}^a = \frac{1}{N} \sum_i A_i \, \phi_i^a \tag{3.14}$$

We have considered the cases $D = 0$ (no feedback of the scalar field on the geometry), $D = 1$ and $D = 2$. For the coefficient of the R^2-term in eq. (2.1) we have taken $a = 0.1$ and $a = 0.001$, motivated by our intention to explore the sensitivity of the results to what is expected to be an irrelevant term. We have considered lattices ranging in size from $8^2 = 64$ to $512^2 = 262144$ sites. The scalar field updates have been performed both by Metropolis Monte Carlo as well as by a heat bath, with compatible results. The number of sweeps for the scalar as well as the gravitational fields varied between at least $500k$ for the 64-site lattice to at least $3.5k$ on the 262144-site lattice. It is only on the largest lattice that we have observed appreciable signs of critical slowing down, leading to somewhat larger errors. The results are shown in Table III and in Figs. 6 ($D = 0$), 7 ($D = 1$) and 8 ($D = 2$).

We have fitted the numerical results for $< \phi^2 >$ to several functional forms, using a standard error-weighted least square algorithm. We estimate the errors in the fits by both computing the intrinsic uncertainty in the fitting parameters, as well as by comparing the variations in the parameters as the number of points used in the fit is changed (we have typically six to seven values of $L = \sqrt{A}$, ranging from 8 to 512, at our disposal). The results clearly suggest a linear behavior of $< \phi^2 >$ in $\ln A$ for all D and a

$$< \phi^2 > \underset{A \to \infty}{\sim} c_0 + c_1 \, \ln A \tag{3.15}$$

but we have tried other fits as well. Assuming the above functional form, we find the results given in Table IV. In the fits typically the χ^2's per degree of freedom are of order one. Note that the coefficients c_0 and c_1 are *not* universal (in particular the coefficient of the $\ln A$ term can be shown to be proportional to the area of an elementary triangle on the lattice, which is clearly not a universal quantity). For the pure $\ln A$ fit there seems to be very little dependence of the fitted coefficients on D.

The above results are not surprising, since without an extrinsic curvature term in the action

$$I_{ec} = \kappa \int d^2x \, \sqrt{g} \, [\, \Delta \phi^a \,]^2 \tag{3.16}$$

where Δ is the covariant Laplacian

$$\Delta = \frac{1}{\sqrt{g}} \, \partial_\mu \, g^{\mu\nu} \, \sqrt{g} \, \partial_\nu \tag{3.17}$$

the surface is expected to completely fold onto itself in embedding space, leading to an infinite fractal dimension, even when the gaussian curvature is zero everywhere ($a \to \infty$)[29,31,32]. If such an embedding in flat euclidean space is considered, then the field ϕ^a is known to play the role of the coordinate $\vec{X}$ in embedding space, as discussed by Polyakov [24]. As far as gravity is concerned, this situation of infinite fractal dimension is of limited consequence, since one does not require the existence of a continuum limit in embedding space, but only for *intrinsic* properties of the manifold. From an intrinsic point of view, the manifold is quite smooth, since $< \phi^2 >$ shows a free-field type behaviour, and the intrinsic or Gaussian curvature (as measured by R^2) is bounded. On the other hand critical properties of 2-d quantum gravity will not necessarily be independent of the existence of an extrinsic curvature term, since correlation functions of ϕ are affected by it[31,32].

If we attempt to fit $< \phi^2 >$ to a power of L instead (as suggested for example by the authors of ref. [28]), namely

$$\ln < \phi^2 > \underset{A \to \infty}{\sim} c_0 + c_1 \ln A \tag{3.18}$$

we find values for $c_1 = 2/d_H$, where d_H is the fractal dimension of the surface, in the range 0.15(2), or d_H about 13. On the other hand the χ^2 parameter for the fits is always at least an order of magnitude larger than in the previously discussed case, and furthermore d_H shows a clear trend towards an increase with lattice size. From this we conclude that the fractal dimension is always infinite in our model. This should be contrasted to the numerical results of the third reference in [14], which gave a finite value for the fractal dimension (but on significantly smaller lattices than the ones we have treated here, and therefore

with understandable uncertainties). Also it is clear that in order to reconcile our results, as well as the semi-classical limit $D \to -\infty$, the power-law term discussed in the second of ref. [28] must have a vanishing coefficient.

We have also investigated the possibility that on the torus and for $D = 1$ there might be a $(\ln A)^2$-term as well, as suggested again by the authors of the second of ref. [28],

$$< \phi^2 > \underset{A \to \infty}{\sim} c_0 + c_1 \ln A + c_2 (\ln A)^2 \tag{3.19}$$

Both for $a = 0.1$ and $a = 0.001$ we obtain the bound

$$c_2 \leq 0.002(2) \tag{3.20}$$

suggesting that if such a term is present, its coefficient must be rather small. On the other hand there is no convincing numerical evidence for such a term in any other model for two-dimensional surfaces. Our results seem to suggest that for the torus the semi-classical ($D \to -\infty$) result is exact, at least for the values of D which we have explored.

It is of interest to explore how some of the above results depend on the gravitational measure. For $D = 1$ and $a = 0$ we have repeated the simulation using the lattice analogue of the DeWitt measure (eq. (3.8)). In the case $D = 1$ this particular measure becomes quite simple, $\prod dl^2$, since all the volume factors cancel out. The computed values for $< \phi^2 >$ are well described by eq. (3.15) (see also Fig. 7), and we find

$$c_0 = 0.19(3) \quad c_1 = 0.175(8) \tag{3.21}$$

which confirms the fact that a change in the measure changes the non-universal coefficients c_0 and c_1, but leaves the functional dependence on A unchanged, and in particular the result $d_H \equiv \infty$.

4. ISING SPINS COUPLED TO GRAVITY

In order to study a model for gravity coupled to Ising spins, we consider the pure gravity action of eq. (2.1), with the additional term

$$\frac{J}{2} \sum_{\text{edges ij}} V_{ij} \left(\frac{S_i - S_j}{l_{ij}} \right)^2 \tag{4.1}$$

which is analogous in form to the scalar field action (3.2), except for the replacement of the scalar field multiplet ϕ_i^a by the Ising spin variables $S_i = \pm 1$.

For the case of the sphere, the Ising model on a dynamically triangulated lattice has been solved exactly by exploiting the equivalence to a large N-matrix model[17]. The critical exponents $\alpha = -1$, $\beta = 1/2$, $\gamma = 2$, $\eta = 2/3$ and $\nu = 3/2$ agree with the conformal field theory estimates of ref. [26]. Here we will investigate again the case of a lattice with the topology of a torus (periodic boundary conditions for all fields), and show that for the torus the semiclassical ($D = -\infty$) results $\alpha = \alpha_0, ...$, where $\alpha_0 = 0, ...$ are the pure Ising critical exponents, appear to hold.

In our simulation we have measured the discrete analogues of the invariant spontaneous magnetization per spin

$$M = \frac{< \int \sqrt{g}\, S >}{< \int \sqrt{g} >} \tag{4.2}$$

and of the zero field susceptibility

$$\chi = \frac{< \int \sqrt{g}\, S \int \sqrt{g}\, S >}{< \int \sqrt{g} >} - \frac{< \int \sqrt{g}\, S >^2}{< \int \sqrt{g} >} \tag{4.3}$$

In a fixed-area ensemble with $A = N = L^2$, these formulae simplify to

$$M = < m > , \qquad m = \frac{1}{A} \sum_i A_i\, S_i \tag{4.4}$$

and

$$\chi = A\,[\,< m^2 > - < m >^2\,] \tag{4.5}$$

respectively. Since on a finite lattice the spontaneous magnetization will vanish identically even at low temperatures, we have found it convenient to also define the quantities M' and χ', which differ from the above expressions by the replacement of m with $|m|$. In addition we have computed the fluctuation in the Ising energy

$$C_{is} = \frac{1}{A}\,[\,< (E_{is})^2 > - < E_{is} >^2\,] \tag{4.6}$$

with

$$E_{is} = \frac{1}{2} \sum_{\text{edges ij}} V_{ij} \left(\frac{S_i - S_j}{l_{ij}}\right)^2 \tag{4.7}$$

If all the edges are taken to be of equal length, then the system reduces to a pure Ising model on a triangular lattice, for which $J_c = \sqrt{3}\ln 3/2 = 0.9514....$ This has provided us with the possibility of a useful comparison of our results with and without gravity on finite lattices. In our simulations we have at first used again the gravitational measure dl/l of eq. (1.8), and have set the higher derivative coupling to $a = 0.001$. We have investigated lattice sizes varying from $8^2 = 64$ sites to $128^2 = 16384$ sites. The length of our runs varies in the critical region ($J_c \approx 1.03$) between 400k sweeps on the smallest lattice and 50k sweeps on the largest lattice.

Let us begin by discussing the magnetization results. Since close to J_c we expect $M \sim (J - J_c)^\beta$, we have that $M^{1/\beta}$ should appear close to linear. In Fig. 9 we show the magnetization M' squared (suggested by the assumption $\beta = 1/2$), and raised to the eight power (suggested by the assumption $\beta = 1/8$), for different size lattices. Our results clearly favor the pure Ising exponent, in spite of the uncertainty in J_c. In the case of the susceptibility we expect from finite size scaling a scaling form

$$\chi(L, J) = L^{2-\eta}\ \bar{\chi}(L^{1/\nu}(J - J_c)) \tag{4.8}$$

Thus in particular the peak in χ (or χ') should scale like $L^{2-\eta}$ for large L. In Table V and in Fig. 10 we show the computed peaks in χ and χ' as a function of L, on a logarithmic scale in the figure. Using a least-square fit we estimate

$$\begin{aligned} 2 - \eta &= 1.78(6) \quad (\text{from } \chi) \\ 2 - \eta &= 1.68(3) \quad (\text{from } \chi') \end{aligned} \tag{4.9}$$

which is completely consistent with the pure Ising results $2 - \eta = \gamma/\nu = 7/4$.

The results for the peak in the Ising specific heat C_{is} as a function of lattice size L are shown again in Table V and in Fig. 11. They resemble closely what is expected from a logarithmic divergence $C_{is} \sim \ln L$ ($\alpha/\nu = 0$). From the combined magnetization, susceptibility and specific heat results, we estimate the critical Ising coupling in the infinite volume limit at

$$J_c = 1.030(5) \tag{4.10}$$

One can improve on the estimate of J_c by considering the fourth-order cumulant[33]

$$U_L(J) = 1 - \frac{< m^4 >}{3 < m^2 >^2} \tag{4.11}$$

with scaling form

$$U_L(J) = \bar{U}(L^{1/\nu}(J - J_c)) \tag{4.12}$$

The curves $U_L(J)$, for different and sufficiently large values of L, should intersect at a common point J_c, where the theory exhibits scale invariance, and where U takes on the fixed point value U^*. We have found that indeed the curves meet very close to a common point, and from the intersection of the curves for $L =$ 8 to 128 we estimate

$$J_c = 1.025(3) \tag{4.13}$$

which is consistent with the previous estimate of the critical point. We also determine $U^* = 0.59(2)$, which is again consistent with the pure Ising model estimate for the invariant charge $U^* \approx 0.58$ [33]. We conclude that for the Ising model coupled to gravity on a simplicial lattice with the topology of a torus, the exponents are, within our errors, the same as in the pure Ising model.

In order to check again the dependence of our results on the specific measure (dl/l) used, we have also performed a simulation with the 'DeWitt' measure appropriate for one field (dl^2), and with the 'Misner' measure $dl^2/\sqrt{V_l}$. We have done the two simulations on a lattice of $32^2 = 1024$ sites, and have chosen $a = 0$. Our conclusions therefore rely mostly on a comparison with similar results with the dl/l measure, and on the same size lattice. While we find that the critical point moves to a larger value ($J_c \approx 1.04$), we find that the peak values in the susceptibility and specific heat are comparable to the values for the dl/l measure on the same size lattice. In Table VI we compare the 'DeWitt' (dl^2) measure for $a = 0$ and $L = 32$ to the 'Misner' ($dl^2/\sqrt{V_l}$) measure for $a = 0$, and to the dl/l-measure results for $a = 0$ and $a = 0.001$. Our results show again little dependence on the measure, at least within our errors and for the lattice sizes considered. A more detailed comparison would involve several lattice sizes, which is outside the scope of the present work. (We did check though that for the 'DeWitt' measure the results on a larger lattice, 64^2, combined with the results on the smaller lattice, are again consistent with pure Ising exponents: for example we estimate $2 - \eta \approx 1.75(14)$). We conclude that the choice of measure does not seem to affect our conclusion that the critical exponents are pure Ising-like.

We have also investigated the critical properties of the Liouville field in the case of gravity coupled to an Ising model. We have computed the Liouville-field susceptibility $\chi_\varphi(J, L)$ discussed previously (in the context of pure gravity) and have found the critical exponent η_φ to be still consistent with 0. In particular for the measure dl/l with $a = 0$ and at the Ising transition, we find using finite size scaling on sizes 8-64 (see Fig. 5) for the exponent $2 - \eta_\varphi = 2.19(19)$, a result which is consistent with the expected free field behavior of a massless Liouville mode. As we move away from the Ising critical point, we still find a similar result, namely that the Liouville susceptibility grows close to linearly in L^2 as L is increased, indicating that the Liouville field behaves like a massless free field for all J.

5. CONCLUSIONS

In the preceding sections we have discussed results relevant for a model of simplicial quantum gravity, and mostly on a manifold withe the topology of a torus. It is characteristic of our model that the variations in the geometry of space are described by fluctuating edge lengths on a lattice with fixed coordination number. We have considered the case of pure gravity ($D = 0$), gravity coupled to a scalar field ($D = 1,\ 2$), and gravity coupled to an Ising model ($D = 1/2$). We have investigated in detail how the results for critical properties depend on what are expected to be irrelevant (R^2-type) terms, as well as on the form of the gravitational measure.

In the case of pure gravity we have computed the exponent γ_χ for both the torus and the sphere, and found good agreement with the expected exact answers from conformal field theory. For the torus we have computed critical properties of the Liouville field, which corresponds to the area density fluctuations on the lattice, and found them to be in agreement with the expectation that the Liouville field behaves like a free massless field for $\lambda \to \lambda_c$, or $A \to \infty$.

By adding a D-component scalar field to the model, we have been able to compute the average $< \phi^2 >$, and therefore make contact with the results on models of random surfaces. These predict, among other things, an infinite fractal dimension for the surface, at least in the absence of extrinsic curvature terms. We found that for the torus the fractal dimension is indeed infinite for $D = 0$, 1 and 2. For $D = 1$ we have looked for an $(\ln A)^2$ term, but have found its amplitude to be quite small and consistent with zero. Our results seem to be insensitive, within the range of parameters explored, to the presence of an R^2-type term in the action or to the detailed form of the measure. We have argued that for sufficiently singular measures though, the triangulation will tend to collapse into a degenerate configuration of edges.

Finally we have studied the case of gravity coupled to an Ising field. By computing the Liouville field susceptibility in the presence of the Ising spins as well (both at the Ising transition and away from it), we have established that the Liouville field stays massless. On the other hand a computation of the

magnetization, susceptibility and specific heat suggests that for the torus the exponents are the same as for the pure Ising model (which is basically the semi-classical $D \rightarrow -\infty$ result). There are two possible explanations for this last finding. The first is that there is some flaw in our model (in the action or in the measure), and that in some other model for lattice gravity coupled to Ising spins the correct gravitational exponents will be obtained. But is appears difficult to reconcile this conclusion with the fact that we seem to obtain the correct value for the pure gravity string susceptibility exponent γ on the sphere, and the fact that the lattice analogue of the Liouville field appears massless (a lack of re-parametrization invariance is expected in general to lead to massive excitations only, with masses of the order of the ultraviolet cutoff). An alternative possible explanation lies with the peculiar properties of the toroidal topology. In the conformal gauge consider the Liouville field, the conformal field theory and the ghosts in a classical background metric $g^0_{\mu\nu}(x)$. Then the trace of the stress-energy tensor is given by[27]

$$< T^{\mu}_{\mu} >= -\frac{C_{tot}}{48\pi} \sqrt{g^0(x)}\, R^0(x)$$

and the requirement $C_{tot} = k + D - 25 = 0$ leads to the $k = 25 - D$-coefficient in front of the effective Liouville action, and to the KPZ results for the scaling dimensions. On the other hand for the torus and for D=0 for example, the only classical solution of the classical equations of motion (derived from the action (2.2) with $\lambda \rightarrow 0$) which is consistent with the boundary conditions is $R^0 = 0$, which would seem to pose no restriction on the value of C_{tot}. We are inclined to conclude therefore that the torus represents possibly a special case, and we expect the Ising exponents to change to the values predicted in refs. [17,26] as we go to the sphere.

In conclusion it is clear that it would be of interest to investigate in greater detail a topology different from the torus, like a sphere or surfaces with boundaries. Even more challenging are of course possible applications to four-dimensional gravity.

ACKNOWLEDGEMENTS

The authors have benefitted from conversations with J. Ambjørn, F. David, V. Kazakov, J. Klauder, P. Menotti, A. Migdal and P. Rakow. A major part of the numerical computations were performed at the San Diego Supercomputer Center. The research of H.W.H. was supported in part by the National Science Foundation under grant NSF-PHY-8906641. The research of M.G. was supported in part by a California State University SCA award.

References

[1] T. Regge, *Nuovo Cimento* **19** (1961) 558.

[2] M. Roček and R. M. Williams, *Phys. Lett.* **104B** (1981) 31 and *Z. Phys.* **C21** (1984) 371.

[3] J. Cheeger, W. Müller and R. Schrader, in the *Heisenberg Symposium*, München 1981, Springer Lecture Notes in Physics, P. Breitlohner and H. P. Dürr eds., New York 1982;
J. Cheeger, W. Müller and R. Schrader, *Comm. Math. Phys.* **92** (1984) 405.

[4] R. Friedberg and T. D. Lee, *Nucl. Phys.* **B242** (1984) 145;
G. Feinberg, R. Friedberg, T. D. Lee and H. C. Ren, *Nucl. Phys.* **B245** (1984) 343.

[5] H. Hamber and R. M. Williams, *Nucl. Phys.* **B248** (1984) 392.
H. Hamber and R. M. Williams, *Nucl. Phys.* **B269** (1986) 712;

[6] H. Hamber, in the proceedings of the *1984 Les Houches Summer School*, Session XLIII, edited by K. Osterwalder and R. Stora, North Holland, 1986.
H. Hamber, in the proceedings of the *1989 Cargèse* NATO workshop on *Probabilistic Methods in Quantum Field Theory and Quantum Gravity*, (Plenum Press, New York 1990).

[7] J. B. Hartle, *J. Math. Phys.* **26** (1985) 804, and **27** (1985) 287;
J. B. Hartle, in the *Proceedings of the Third Moskow Seminar on Quantum Gravity*, M. Markov, V. Berezin and V. Frolov eds., World Scientific 1985;
J. B. Hartle, *Class. Quantum Grav.* **2** (1985) 707.
[8] H. Hamber and R. M. Williams, *Phys. Lett.* **157B** (1985) 368,
[9] H. Hamber and R. M. Williams, *Nucl. Phys.* **B267** (1986) 482.
[10] B. Berg, *Phys. Rev. Lett.* **55** (1985) 904 and *Phys. Lett.* **176B** (1986) 39.
[11] M. Bander and C. Itzykson, *Nucl. Phys.* **B257 [FS14]** (1985) 531;
A. Jevicki and M. Ninomiya, *Phys. Lett.* **150B** (1985) 115 and *Phys. Rev.* **D33** (1986) 1634;
Z. Hlousek and K. O'Brien, Cornell preprint CNLS 88/860 (1988) (unpublished).
[12] D. A. Eliezer, *Nucl. Phys.* **B319** (1989) 667.
[13] F. David, *Nucl. Phys.* **B257 [FS14]** (1985) 45 and 543;
A. Billoire and F. David, *Nucl. Phys.* **B275** (1986) 617;
F. David, J. Jurkiewicz, A. Krzywicki and B. Petersen, *Nucl. Phys.* **B290** (1987) 218;
[14] V. A. Kazakov, *Phys. Lett.* **150B** (1985) 282;
V. A. Kazakov, I. K. Kostov and A. A. Migdal, *Phys. Lett.* **157B** (1985) 295;
D. V. Boulatov, V. A. Kazakov, I. K. Kostov and A. A. Migdal, *Nucl. Phys.* **B275** (1986) 641;
D. V. Boulatov and V. A. Kazakov, *Phys. Lett.* **186B** (1987) 379;
V. A. Kazakov and A. A. Migdal, *Nucl. Phys.* **B311** (1988) 171.
M. Agishtein and A. A. Migdal, Princeton preprint PUPT-1171 (1990).
[15] J. Ambjørn, B. Durhuus and J. Frölich, *Nucl. Phys.* **B257 [FS14]** (1985) 433;
[16] E. Brezin and V. A. Kazakov, *Phys. Lett.* **119A** (1986) 140;
M. Douglas and S. Shenker, *Nucl. Phys.* **B335** (1989) 635;
D. Gross and A. Migdal, *Phys. Rev. Lett.* **64** (1990) 127;
G. Parisi, Rome preprint 89/30 (1989).
[17] V. A. Kazakov, *Phys. Lett.* **119A** (1986) 140;
D. V. Boulatov and V. A. Kazakov, *Phys. Lett.* **186B** (1987) 379.
[18] A. Das, M. Kaku and P. K. Townsend, *Phys. Lett.* **81B** (1979) 11;
L. Smolin, *Nucl. Phys.* **B148** (1979) 333;
C. Mannion and J. G. Taylor, *Phys. Lett.* **100B** (1981) 261;
M. Kaku, *Phys. Rev.* **D27** (1983) 2819;
E. T. Tomboulis, *Phys. Rev. Lett.* **52** (1984) 1173;
K. Kondo, *Prog. Theor. Phys.* **72** (1984) 841.
[19] P. Menotti and A. Pelissetto, *Ann. Phys. (NY)* **170** (1986) 287, *Nucl. Phys.* **B288** (1987) 813 and *Phys. Rev.* **D35** (1987) 1194;
S. Caracciolo and A. Pelissetto, *Nucl. Phys.* **B299** (1988) 693, and *Phys. Lett.* **207B** (1988) 468.
[20] N. P. Konopleva and V. N. Popov, *'Gauge Fields'*, Harwood Academic Publishers, New York 1979, and references therein.
[21] B. DeWitt, in *'General Relativity - An Einstein centenary survey'*, edited by S.W. Hawking and W. Israel, Cambridge University press (1979), and references therein;
K. Fujikawa, *Nucl. Phys.* **B226** (1983) 437.
[22] H. Leutwyler, *Phys. Rev.* **134** (1964) 1155;
E. S. Fradkin and G. A. Vilkovisky, *Phys. Rev.* **D8** (1973) 4241;
E. S. Fradkin and G. A. Vilkovisky, CERN preprint TH-2332 (1977), unpublished.
[23] V. De Alfaro, S. Fubini and G. Furlan, *Nuovo Cimento* **57B** (1980) 227 and **76A** (1983) 365;
P. Menotti and A. Pelissetto, *Phys. Rev.* **D35** (1987) 1194;
M. Bander, *Phys. Rev. Lett.* **57** (1986) 1825.
[24] A. M. Polyakov, *Phys. Lett.* **103B** (1981) 207.
[25] A. M. Polyakov, *Mod. Phys. Lett.* **A2** (1987) 899.
[26] V. G. Knizhnik, A. M. Polyakov and A. B. Zamolodchikov, *Mod. Phys. Lett.* **A3** (1988) 819.
[27] F. David, *Mod. Phys. Lett.* **A3** (1988) 1651; 207;
F. David and E Guitter, *Europhys. Lett.* **3** (1987) 1169, and *Nucl. Phys.* **B293** (1988) 332.
[28] J. Distler and H. Kawai, *Nucl. Phys.* **B321** (1989) 509;

J. Distler, Z. Hlousek and H. Kawai, *Int. J. Mod. Phys.* **A5** (1990) 1093.
[29] D. Gross, *Phys. Lett.* **138B** (1984) 185;
A. Billoire, D. Gross and E. Marinari, *Phys. Lett.* **139B** (1984) 75.
[30] N. Christ, R. Friedberg and T. D. Lee, *Nucl. Phys.* **B202** (1982) 89 and *Nucl. Phys.* **B210 [FS6]** (1982) 310,337;
C. Itzykson, *Cargèse Lecture Notes 1983*, published in the proceedings (Plenum Press, New York, 1985).
[31] W. Helfrich, *J. Phys.* **46** (1985) 1263;
L. Peliti and S. Leibler, *Phys. Rev. Lett.* **54** (1985) 1690;
D. Foerster, *Phys. Lett.* **A114** (1986) 115;
H. Kleinert, *Phys. Lett.* **B174** (1986) 335;
A. M. Polyakov, *Nucl. Phys.* **B268** (1986) 406.
F. David and E Guitter, *Nucl. Phys.* **B295** (1988) 332;
[32] Y. Kantor and D. Nelson, *Phys. Rev. Lett.* **58** (1987) 2774;
J. Ambjørn, B. Durhuus and T. Jonsson, Niels Bohr Institute preprint NBI-HE-88-61;
M. Baig, D. Espriu and J. F. Wheater, *Nucl. Phys.* **B** *Proc. Suppl.* **4** (1988) 88;
J. Kogut, A. Moreo and S. Das, *Nucl. Phys.* **B327** (1989) 530;
S. Catterall, *Phys. Lett.* **B220** (1989) 207;
R. Renken and J. B. Kogut, Illinois preprints 90-5 and 90-14 (1990);
C. Baillie, D. Johnston and R. Williams, Caltech preprints (1990).
[33] K. Binder, *Phys. Rev. Lett.* **47** (1981) 693;
K. Binder, *Z. Phys.* **B43** (1981) 119.

Table captions

Table I: Average of the curvature squared as a function of the total area for the torus and the sphere

Table II: Liouville field or area density susceptibility as a function of lattice size. $D = 0$ refers to pure gravity, and $D = 1/2$ to gravity coupled to an Ising model.

Table III: Expectation values $< \phi^2 >$ as a function of lattice size, for different values of a and D. The column labelled with a $*$ corresponds to the case of the DeWitt (dl^2) measure.

Table IV: Summary of least square fits for the coefficients c_0 and c_1 in eq. (3.15) for $< \phi^2 >$, for different values of D and a.

Table V: Peak values for the Ising susceptibilities χ and χ', and for the specific heat C, as a function of lattice size.

Table VI: Results for the peak values of the Ising susceptibilities χ and χ', and of the specific heat C, on a 32^2-lattice and for several forms of the gravitational measure.

Table I.

	$<\sum_i \delta_i^2/A_i>/A$	
A	torus	sphere
16	0.40977(19)	1.10221(77)
25	0.41155(17)	0.74625(138)
36	0.41088(19)	0.60633(200)
64	0.40968(8)	0.48921(59)
100	0.40963(8)	0.44976(10)
144	0.40982(9)	0.43582(38)
256	0.40983(9)	0.42755(26)
1024	0.41010(9)	0.42349(26)
4096	0.40995(10)	0.41818(40)

Table II.

	$\chi_\phi(L)$		
L	D=0	D=1/2 (J_c)	D=1/2 (all J)
8	1.58(18)	1.8(2)	1.57(18)
16	10.18(3.2)	7.2(5)	9.9(3.2)
32	49.3(14.)	22.5(4.)	30.0(16.2)
64	113.87(33.)	211.(50.)	129.(60.)
128	409.19(220.)	60.(70.)	83.(51.)

Table III.

$A=L^2$	$<\phi^2>$					
	D=0		D=1			D=2
	a=0.1	a=0.001	a=0.1	a=0.001	a=0*	a=0.1
64	0.902(8)	0.915(6)	0.919(1)	0.949(1)	0.916(4)	0.938(1)
256	1.164(6)	1.196(6)	1.186(1)	1.214(2)	1.162(4)	1.210(1)
1024	1.420(4)	1.455(6)	1.443(3)	1.487(4)	1.402(4)	1.478(3)
4096	1.698(21)	1.702(12)	1.702(6)	1.743(9)	1.646(9)	1.734(6)
16384	1.928(47)	1.984(35)	1.919(17)	1.923(22)	1.906(30)	2.012(17)
65536	2.321(161)	2.117(39)	2.340(65)	2.165(47)	2.212(70)	2.191(65)
262144	2.44(23)	2.50(13)	2.327(27)	2.424((32)	2.36(8)	-

Table IV.

D	a	c_0	c_1
0	0.1	0.13(3)	0.187(3)
0	0.001	0.17(3)	0.185(3)
1	0.1	0.15(3)	0.187(3)
1	0.001	0.16(3)	0.190(3)
2	0.1	0.14(3)	0.193(3)

Table V.

L	$\chi_{max}(L)$	$\chi'_{max}(L)$	$C_{max}(L)$
8	24.6(10)	4.5(2)	4.16(5)
16	97.3(50)	14.5(5)	4.89(5)
32	236.(60.)	51.(4.)	5.74(10)
64	710.(120.)	179.(50.)	6.62(60)
128	359.(150.)	342.(70.)	6.01(1.2)

Table VI.

Measure	χ_{max}	χ'_{max}	C_{max}
DW, a=0	279(40)	49(5)	6.4(4)
M, a=0	332(40)	51(6)	6.8(6)
dl/l, a=0	353(51)	37(6)	5.5(6)
dl/l, a=0.001	331(55)	40(6)	5.1(5)

Figure captions

Figure 1: Two equivalent triangulation of a smooth manifold, using a lattice of coordination number six.

Figures 2,3: Average curvature squared as a function of the lattice area, for $a = 1.0$. The two graphs correspond to the torus (2) and the sphere (3).

Figure 4: Distribution of curvatures $\delta_i \sim \sqrt{g}\, R$ on a lattice with $512^2 = 262144$ sites.

Figure 5: Area fluctuation or Liouville susceptibility, as a function of lattice size. The circles correspond to pure gravity and $a = 0$ ($D = 0$), and the squares to gravity with $a = 0$ coupled to an Ising field ($D = 1/2$). The straight line on the logarithmic scale indicates a growth proportional to L^2.

Figure 6: Scalar field average $< \phi^2 >$ for pure gravity ($D = 0$), with $a = 0.1$ (circles) and $a = 0.001$ (squares). The straight line corresponds to a logarithmic divergence, or $d_H = \infty$.

Figure 7: Scalar field average $< \phi^2 >$ for gravity coupled to a massless scalar field ($D = 1$), with $a = 0.1$ (circles) and $a = 0.001$ (squares). The diamonds indicate results for the DeWitt measure (dl^2), with $a = 0$. The straight line corresponds to a logarithmic divergence, or $d_H = \infty$.

Figure 8: Scalar field average $< \phi^2 >$ for gravity coupled to a two-component massless scalar field ($D = 2$), with $a = 0.1$. The straight line corresponds to a logarithmic divergence, or $d_H = \infty$.

Figure 9: Ising magnetization M' squared and raised to the eight power, for different lattice sizes ($L =$ 8 (diagonal crosses), 16 (crosses), 32 (squares), 64 (circles), 128 (diamonds)). The pure Ising magnetization on an $L = 64$ lattice, raised to the eight power, (diamonds) is shown for comparison.

Figure 10: Peak values for the Ising susceptibilities χ (circles) and χ' (diamonds) for different size lattices. The straight lines correspond to an exponent $2 - \eta = 7/4$.

Figure 11: Peak values for the Ising specific heat C. The straight line corresponds to a logarithmic divergence ($\alpha = 0$).



Figure 1



Figure 2



Figure 3



Figure 4



Figure 5



Figure 6



Figure 7



Figure 8



Figure 9



Figure 10



Figure 11